

# TWISTED $\gamma$ -FILTRATION OF A LINEAR ALGEBRAIC GROUP.

KIRILL ZAINOULLINE

ABSTRACT. In the present notes we introduce and study the twisted  $\gamma$ -filtration on  $K_0(G_s)$ , where  $G_s$  is a split simple linear algebraic group over a field  $k$  of characteristic prime to the order of the center of  $G_s$ . We apply this filtration to construct torsion elements in the  $\gamma$ -ring of the variety of complete  $G$ -flags, where  $G$  is an inner form of  $G_s$ .

## 1. INTRODUCTION

Let  $X$  be a smooth projective variety over a field  $k$ . Consider the Grothendieck  $\gamma$ -filtration on  $K_0(X)$ . It is given by the ideals [6, §2.3] (see also [8, §2])

$$\gamma^i K_0(X) = \langle c_{n_1}(b_1) \cdots c_{n_m}(b_m) \mid n_1 + \dots + n_m \geq i, b_1, \dots, b_m \in K_0(X) \rangle, i \geq 0$$

generated by products of Chern classes in  $K_0$ . Let  $\gamma^i(X)$  denote the  $i$ -th subsequent quotient and let  $\gamma^*(X) = \bigoplus_{i \geq 0} \gamma^i(X)$  denote the associated graded commutative ring called the  $\gamma$ -ring of  $X$ .

The ring  $\gamma^*(X)$  was invented by Grothendieck to approximate the topological filtration on  $K_0$  and, hence, the Chow ring  $\mathrm{CH}^*(X)$  of algebraic cycles modulo rational equivalence. Indeed, by the Riemann-Roch theorem (see [6, §2]) the  $i$ -th Chern class  $c_i$  induces an isomorphism with  $\mathbb{Q}$ -coefficients, i.e.  $c_i: \gamma^i(X; \mathbb{Q}) \xrightarrow{\sim} \mathrm{CH}^i(X; \mathbb{Q})$ . Moreover, in some cases the ring  $\gamma^*(X)$  can be used to compute  $\mathrm{CH}^*(X)$ , e.g.  $\gamma^1(X) = \mathrm{CH}^1(X)$  and there is a surjection  $\gamma^2(X) \twoheadrightarrow \mathrm{CH}^2(X)$  (see [7, Ex. 15.3.6]).

In the present notes we provide a uniform lower bound for the torsion part of  $\gamma^*(X)$ , where  $X = {}_\xi \mathfrak{B}_s$  is a twisted form of the variety of Borel subgroups  $\mathfrak{B}_s$  of a split simple linear algebraic group  $G_s$  by means of a cocycle  $\xi \in H^1(k, G_s)$ . Note that the groups  $\gamma^2(X)$  and  $\gamma^3(X)$  have been studied for  $G_s = \mathrm{PGL}_n$  in [8] and for strongly inner forms in [4]. In particular, it was shown in [4, §3,7] that in the strongly inner case the torsion part of  $\gamma^2(X)$  determines the Rost invariant.

Our main tool is the twisted  $\gamma$ -filtration on  $K_0(G_s)$ , where  $G_s$  is a split simple linear algebraic group. Roughly speaking, it is defined to be the image (see Definition 4.4) of the  $\gamma$ -filtration on  $K_0$  of the twisted form  $X$  under the composition  $K_0(X) \rightarrow K_0(\mathfrak{B}_s) \rightarrow K_0(G_s)$ , where the first map is given by the restriction and the second map is induced by taking the quotient.

Let  $\gamma_\xi^*$  denotes the associated graded ring of the twisted  $\gamma$ -filtration. It has the following important properties:

- (i) The ring  $\gamma_\xi^*$  can be explicitly computed (see Theorem 4.5). Observe that  $\gamma_\xi^0 = \mathbb{Z}$ ,  $\gamma_\xi^1 = 0$  and  $\gamma_\xi^i$  is torsion for  $i > 1$ .
- (ii) There is a surjective ring homomorphism  $\gamma^*(X) \twoheadrightarrow \gamma_\xi^*$ . Hence,  $\gamma_\xi^*$  gives a lower bound for the  $\gamma$ -ring of the twisted form  $X = {}_\xi \mathfrak{B}_s$ .

- (iii) The assignment  $\xi \mapsto \gamma_\xi^*$  respects the base change and, therefore, is an invariant of a  $G_s$ -torsor  $\xi$ , moreover, the ring  $\gamma_\xi^*$  can be viewed as a substitute for the  $\gamma$ -ring of the inner group  ${}_\xi G_s$ .

In the last section we use these properties to construct nontrivial torsion elements in  $\gamma^2(X)$  for some twisted flag varieties  $X$  (see 5.3 and 5.5). In particular, we establish the connection between the indexes of the Tits algebras of  $\xi$  and the order of the special cycle  $\theta \in \gamma^2(X)$  constructed in [4].

## 2. PRELIMINARIES.

In the present section we recall several basic facts concerning linear algebraic groups, characters and the Grothendieck  $K_0$  (see [9, §24], [4, §1B, §6]).

2.1. Let  $G_s$  be a split simple linear algebraic group of rank  $n$  over a field  $k$ . We assume that characteristic of  $k$  is prime to the order of the center of  $G_s$ . We fix a split maximal torus  $T$  and a Borel subgroup  $B$  such that  $T \subset B \subset G_s$ .

Let  $\Lambda_r$  and  $\Lambda$  be the root and the weight lattices of the root system of  $G_s$  with respect to  $T \subset B$ . Let  $\{\alpha_1, \dots, \alpha_n\}$  be a set of simple roots (a basis of  $\Lambda_r$ ) and let  $\{\omega_1, \dots, \omega_n\}$  be the respective set of fundamental weights (a basis of  $\Lambda$ ), i.e.  $\alpha_i^\vee(\omega_j) = \delta_{ij}$ . The group of characters  $T^*$  of  $T$  is an intermediate lattice  $\Lambda_r \subset T^* \subset \Lambda$  that determines the isogeny class of  $G_s$ . If  $T^* = \Lambda$ , then the group  $G_s$  is simply connected and if  $T^* = \Lambda_r$  it is adjoint.

2.2. Let  $\mathbb{Z}[T^*]$  be the integral group ring of  $T^*$ . Its elements are finite linear combinations  $\sum_i a_i e^{\lambda_i}$ ,  $\lambda_i \in T^*$ . Let  $\mathfrak{B}_s$  denote the variety of Borel subgroups  $G_s/B$  of  $G_s$ . Consider the characteristic map for  $K_0$  (see [3, §2.8])

$$\mathbf{c}: \mathbb{Z}[T^*] \rightarrow K_0(\mathfrak{B}_s)$$

defined by sending  $e^\lambda$ ,  $\lambda \in T^*$ , to the class of the associated line bundle  $[\mathcal{L}(\lambda)]$ . Observe that the ring  $K_0(\mathfrak{B}_s)$  does not depend on the isogeny class of  $G_s$  while the group of characters  $T^*$  and, hence, the image of  $\mathbf{c}$  does.

Since  $K_0(\mathfrak{B}_s)$  is generated by the classes  $[\mathcal{L}(\omega_i)]$ ,  $i = 1 \dots n$ , the characteristic map  $\mathbf{c}$  is surjective if  $G_s$  is simply connected. If  $G_s$  is adjoint, then the image of  $\mathbf{c}$  is generated by the classes  $[\mathcal{L}(\alpha_i)]$ , where

$$\alpha_i = \sum_j c_{ij} \omega_j \quad \text{and, therefore,} \quad \mathcal{L}(\alpha_i) = \otimes_j \mathcal{L}(\omega_j)^{\otimes c_{ij}},$$

and  $c_{ij} = \alpha_i^\vee(\alpha_j)$  are the coefficients of the Cartan matrix of  $G_s$ .

2.3. The Weyl group  $W$  of  $G_s$  acts on weights via simple reflections  $s_{\alpha_i}$  as

$$s_{\alpha_i}(\lambda) = \lambda - \alpha_i^\vee(\lambda) \alpha_i, \quad \lambda \in \Lambda.$$

For each element  $w \in W$  we define (cf. [13, §2.1]) the weight  $\rho_w \in \Lambda$  as

$$\rho_w = \sum_{\{i \in 1 \dots n \mid w^{-1}(\alpha_i) < 0\}} w^{-1}(\omega_i).$$

In particular, for a simple reflection  $w = s_{\alpha_j}$  we have

$$\rho_w = \sum_{\{i \in 1 \dots n \mid s_{\alpha_j}(\alpha_i) < 0\}} s_{\alpha_j}(\omega_i) = s_{\alpha_j}(\omega_j) = \omega_j - \alpha_j.$$

Observe that the quotient  $\Lambda/\Lambda_r$  coincides with the group of characters of the center of the simply connected cover of  $G_s$ . Since  $W$  acts trivially on  $\Lambda/\Lambda_r$ , we have

$$\bar{\rho}_w = \sum_{\{i \in 1 \dots n \mid w^{-1}(\alpha_i) < 0\}} \bar{\omega}_i \in \Lambda/T^*,$$

where  $\bar{\rho}_w$  denotes the class of  $\rho_w \in \Lambda$  modulo  $T^*$ . In particular,  $\bar{\omega}_i = \bar{\rho}_{s_{\alpha_i}}$ .

2.4. Let  $\mathbb{Z}[\Lambda]^W$  denote the subring of  $W$ -invariant elements. Then the integral group ring  $\mathbb{Z}[\Lambda]$  is a free  $\mathbb{Z}[\Lambda]^W$ -module with the basis  $\{e^{\rho_w}\}_{w \in W}$  (see [13, Thm.2.2]). Now let  $\epsilon: \mathbb{Z}[\Lambda] \rightarrow \mathbb{Z}$ ,  $e^\lambda \mapsto 1$  be the augmentation map. By the Chevalley Theorem the kernel of the surjection  $\mathfrak{c}$  is generated by elements  $x \in \mathbb{Z}[\Lambda]^W$  such that  $\epsilon(x) = 0$ . Hence, there is an isomorphism

$$\mathbb{Z}[\Lambda] \otimes_{\mathbb{Z}[\Lambda]^W} \mathbb{Z} \simeq \mathbb{Z}[\Lambda] / \ker(\mathfrak{c}) \simeq K_0(\mathfrak{B}_s).$$

So the elements

$$\{g_w = \mathfrak{c}(e^{\rho_w}) = [\mathcal{L}(\rho_w)]\}_{w \in W}$$

form a  $\mathbb{Z}$ -basis of  $K_0(\mathfrak{B}_s)$  called the Steinberg basis.

2.5. Following [14] we associate with each  $\chi \in \Lambda/T^*$  and each cocycle  $\xi \in Z^1(k, G_s)$  the central simple algebra  $A_{\chi, \xi}$  over  $k$  called the Tits algebra. This defines a group homomorphism

$$\beta_\xi: \Lambda/T^* \rightarrow Br(k) \text{ with } \beta_\xi(\chi) = [A_{\chi, \xi}].$$

Let  $\mathfrak{B} = {}_\xi \mathfrak{B}_s$  denote the twisted form of the variety of Borel subgroups  $\mathfrak{B}_s$  by means of  $\xi$ . Consider the restriction map on  $K_0$  over the separable closure  $k_{sep}$

$$res: K_0(\mathfrak{B}) \rightarrow K_0(\mathfrak{B} \times_k k_{sep}) = K_0(\mathfrak{B}_s),$$

where we identify  $K_0(\mathfrak{B} \times_k k_{sep})$  with  $K_0(\mathfrak{B}_s)$ . By [11, Thm.4.2] the image of the restriction can be identified with the sublattice

$$\langle \iota_w \cdot g_w \rangle_{w \in W},$$

where  $g_w = [\mathcal{L}(\rho_w)]$  is an element of the Steinberg basis and  $\iota_w = \text{ind}(\beta_\xi(\bar{\rho}_w))$  is the index of the respective Tits algebra. Observe that if  $G_s$  is simply connected, then all indexes  $\iota_w$  are trivial and the restriction map becomes an isomorphism.

### 3. THE $K_0$ OF A SPLIT SIMPLE (ADJOINT) GROUP

In the present section we provide an explicit description of the ring  $K_0(G_s)$  in terms of generators and relations for every simple split linear algebraic group  $G_s$ . The method to compute  $K_0(G_s)$  was known before, however, due to the lack of precise references we provide the computations here.

3.1. **Definition.** Let  $\mathfrak{c}: \mathbb{Z}[\Lambda] \rightarrow K_0(\mathfrak{B}_s)$  be the characteristic map for the simply connected cover of  $G_s$ . We define the ring  $\mathfrak{G}_s$  to be the quotient

$$\mathfrak{G}_s := \mathbb{Z}[\Lambda/T^*] / \overline{(\ker \mathfrak{c})}$$

and the surjective ring homomorphism  $q$  to be the composite

$$q: K_0(\mathfrak{B}_s) \xrightarrow[\simeq]{\mathfrak{c}^{-1}} \mathbb{Z}[\Lambda] / (\ker \mathfrak{c}) \twoheadrightarrow \mathbb{Z}[\Lambda/T^*] / \overline{(\ker \mathfrak{c})} = \mathfrak{G}_s.$$

Observe that if  $G_s$  is simply connected, then  $\mathfrak{G}_s = \mathbb{Z}$ .

**3.2. Remark.** By [10, Cor.33] applied to  $X = G_s$  and to the simply-connected cover  $G = \hat{G}_s$  of  $G_s$ , there is an isomorphism

$$K_0(G_s) \simeq \mathbb{Z} \otimes_{R(\hat{G}_s)} K_0(\hat{G}_s, G_s),$$

where  $R(\hat{G}_s) \simeq \mathbb{Z}[\Lambda]^W$  is the representation ring. By [10, Cor.5] applied to  $G = \hat{G}_s$ ,  $X = \text{Spec } k$  and  $G/H = G_s$  there is an isomorphism

$$K_0(\hat{G}_s, G_s) \simeq R(H),$$

where  $R(H) \simeq \mathbb{Z}[\Lambda/T^*]$  is the representation ring. Therefore,

$$K_0(G_s) \simeq \mathbb{Z} \otimes_{\mathbb{Z}[\Lambda]^W} \mathbb{Z}[\Lambda/T^*] \simeq \mathfrak{G}_s.$$

**3.3. Lemma.** *The ideal  $\overline{(\ker \mathfrak{c})} \subset \mathbb{Z}[\Lambda/T^*]$  is generated by the elements*

$$d_i(1 - e^{\bar{\omega}_i}), \quad i = 1 \dots n,$$

where  $d_i$  is the dimension of the  $i$ -th fundamental representation.

*Proof.* By the Chevalley Theorem the subring of invariants  $\mathbb{Z}[\Lambda]^W$  can be identified with the polynomial ring  $\mathbb{Z}[\rho_1, \dots, \rho_n]$ , where  $\rho_i$  is the  $i$ -th fundamental representation, i.e.

$$\rho_i = \sum_{\lambda \in W(\omega_i)} e^\lambda$$

(here  $W(\omega_i)$  denotes the  $W$ -orbit of the fundamental weight  $\omega_i$ ).

Since  $d_i = \epsilon(\rho_i)$ ,  $\ker \mathfrak{c} = (d_1 - \rho_1, \dots, d_n - \rho_n)$ . To finish the proof observe that  $\overline{(d_i - \rho_i)} = d_i(1 - e^{\bar{\omega}_i})$ .  $\square$

**3.4. Remark.** Observe that by definition and 3.3 we have  $\mathfrak{G}_s \otimes \mathbb{Q} \simeq \mathbb{Q}$ .

**3.5.** In the following examples we compute the ring  $\mathfrak{G}_s \simeq K_0(G_s)$  for every simple split linear algebraic group  $G_s$  (we refer to [9, §24] for the description of  $\Lambda/T^*$  and to [1, Ch.8, Table 2] for the dimensions of fundamental representations).

$\Lambda/T^*$	$G_s, m \geq 1$	Example
$\mathbb{Z}/m\mathbb{Z}, m \geq 2$	$SL_{n+1}/\mu_m$	(3.6)
$\mathbb{Z}/2\mathbb{Z}$	$O_{m+4}^+, PSp_{2m+2}, HSpin_{4m+4}, E_7^{ad}$	(3.7)
$\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$	$P GO_{4m+4}^+$	(3.8)
$\mathbb{Z}/3\mathbb{Z}$	$E_6^{ad}$	(3.9)
$\mathbb{Z}/4\mathbb{Z}$	$P GO_{4m+2}^+$	(3.10)

**3.6. Example.** Consider the case  $G_s = SL_{n+1}/\mu_m, m \geq 2$ . The group  $G_s$  has type  $A_n$  and  $\Lambda/T^* = \langle \sigma \rangle$  is cyclic of order  $m$ . The quotient map  $\Lambda/\Lambda_r \rightarrow \Lambda/T^*$  sends  $\bar{\omega}_i \in \Lambda/\Lambda_r, i = 1 \dots n$  to  $(i \bmod m)\sigma \in \Lambda/T^*$ .

By Definition 3.1 and Lemma 3.3 we have

$$\mathfrak{G}_s \simeq \mathbb{Z}[y]/(1 - (1 - y)^m, d_1 y, \dots, d_{m-1} y^{m-1}),$$

where  $y = (1 - e^\sigma)$  and  $d_j = \gcd\{\binom{n+1}{i} \mid i \equiv j \bmod m, i = 1 \dots n\}$ .

In particular, for  $G_s = SL_p/\mu_p = PGL_p$ , where  $p$  is a prime, we obtain

$$\mathfrak{G}_s \simeq \mathbb{Z}[y]/(\binom{p}{1}y, \binom{p}{2}y^2, \dots, \binom{p}{p-1}y^{p-1}, y^p).$$

**3.7. Example.** Assume that  $\Lambda/T^* = \langle \sigma \rangle$  has order 2. Then

$$\mathfrak{G}_s \simeq \mathbb{Z}[y]/(y^2 - 2y, dy),$$

where  $y = (1 - e^\sigma)$  and  $d$  is the g.c.d. of dimensions of representations corresponding to  $\omega_i$  with  $\bar{\omega}_i = \sigma$ . The integer  $d$  can be determined as follows:

$B_n$ : We have  $\Lambda/\Lambda_r = \{0, \bar{\omega}_n\} \simeq \mathbb{Z}/2\mathbb{Z}$  which corresponds to the adjoint group  $G_s = O_{2n+1}^+$ . Since  $\bar{\omega}_i = 0$  for each  $i \neq n$ ,  $d$  coincides with the dimension of  $\omega_n$  that is  $2^n$ .

$C_n$ : We have  $\Lambda/\Lambda_r = \{0, \sigma = \bar{\omega}_1 = \bar{\omega}_3 = \dots\} \simeq \mathbb{Z}/2\mathbb{Z}$  that is  $G_s = PSp_{2n}$ . Since  $\bar{\omega}_i = 0$  for even  $i$ ,  $d$  is the g.c.d. of dimensions of  $\omega_1, \omega_3, \dots$ , i.e.

$$d = \gcd(2n, \binom{2n}{3} - \binom{2n}{1}, \binom{2n}{5} - \binom{2n}{3}, \dots).$$

$D_n$ : If  $n$  is odd, then  $\Lambda/\Lambda_r = \{0, \bar{\omega}_{n-1}, \bar{\omega}_1, \bar{\omega}_n\} \simeq \mathbb{Z}/4\mathbb{Z}$ , where  $\bar{\omega}_1 = 2\bar{\omega}_{n-1} = 2\bar{\omega}_n$ . Therefore,  $\Lambda/T^* \simeq \mathbb{Z}/2\mathbb{Z}$  if it is a quotient of  $\Lambda/\Lambda_r$  modulo the subgroup  $\{0, \bar{\omega}_1\}$ . In this case  $\Lambda/T^* = \{0, \sigma = \bar{\omega}_{n-1} = \bar{\omega}_n\}$  which corresponds to the special orthogonal group  $G_s = O_{2n}^+$ . Since  $\bar{\omega}_s = s\bar{\omega}_1$  for  $2 \leq s \leq n-2$  and  $\bar{\omega}_1 = 0$  in  $\Lambda/T^*$ ,  $d$  is the g.c.d. of the dimensions of  $\omega_{n-1}$  and  $\omega_n$  that is  $2^{n-1}$ .

If  $n$  is even, then  $\Lambda/\Lambda_r = \{0, \bar{\omega}_{n-1}\} \oplus \{0, \bar{\omega}_n\} \simeq \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$ , where  $\bar{\omega}_1 = \bar{\omega}_{n-1} + \bar{\omega}_n$ . In this case, we have two cases for  $\Lambda/T^*$ :

- (1) It is the quotient of  $\Lambda/\Lambda_r$  modulo the diagonal subgroup  $\{0, \bar{\omega}_{n-1} + \bar{\omega}_n\}$ . Then  $\Lambda/T^* = \{0, \sigma = \bar{\omega}_{n-1} = \bar{\omega}_n\}$ ,  $G_s = O_{2n}^+$  and  $d$  is the same as in the odd case, i.e.  $d = 2^{n-1}$ .
- (2) It is the quotient modulo one of the factors, e.g.  $\Lambda/T^* = \{0, \sigma = \bar{\omega}_{n-1}\}$ , where  $\bar{\omega}_n = 0$ . This corresponds to the half-spin group  $G_s = HSpin_{2n}$ . We have  $\bar{\omega}_1 = \bar{\omega}_3 = \dots = \bar{\omega}_{n-1}$  and  $\bar{\omega}_i = 0$  if  $i$  is even. Therefore,  $d = \gcd(2n, \binom{2n}{3}, \dots, \binom{2n}{n-3}, 2^{n-1})$  which implies that  $d = 2^{v_2(n)+1}$ , where  $v_2(n)$  denotes the 2-adic valuation of  $n$ .

$E_7$ : We have  $\Lambda/\Lambda_r = \{0, \sigma = \bar{\omega}_7 = \bar{\omega}_5 = \bar{\omega}_2\} \simeq \mathbb{Z}/2\mathbb{Z}$  with  $\bar{\omega}_1 = \bar{\omega}_3 = \bar{\omega}_4 = \bar{\omega}_6 = 0$ . Therefore,  $d = \gcd(56, \binom{56}{3}, 912) = 8$ .

**3.8. Example.** Assume that  $\Lambda/T^* = \langle \sigma_1 \rangle \oplus \langle \sigma_2 \rangle$ , where  $\sigma_1$  and  $\sigma_2$  are of order 2. In this case  $G_s = PGO_{2n}^+$  is an adjoint group ( $T^* = \Lambda_r$ ) of type  $D_n$  with  $n$  even. We have  $\sigma_1 = \bar{\omega}_{n-1}$  and  $\sigma_2 = \bar{\omega}_n$ ,  $\bar{\omega}_s = s\bar{\omega}_1$ ,  $2 \leq s \leq n-2$ ,  $2\bar{\omega}_1 = 0$  and  $\bar{\omega}_1 = \bar{\omega}_{n-1} + \bar{\omega}_n$ . Then

$$\mathfrak{G}_s \simeq \mathbb{Z}[y_1, y_2]/(y_1^2 - 2y_1, y_2^2 - 2y_2, d_1y_1, d_2y_2, d(y_1 + y_2 - y_1y_2)),$$

where  $y_1 = (1 - e^{\sigma_1})$ ,  $y_2 = (1 - e^{\sigma_2})$ ;  $d_1$  (resp.  $d_2$ ) is the g.c.d. of dimensions of  $\omega_i$  with  $\bar{\omega}_i = \bar{\omega}_{n-1}$  (resp.  $\bar{\omega}_i = \bar{\omega}_n$ ) that is  $d_1 = d_2 = 2^{n-1}$ ; and  $d$  is the g.c.d. of dimensions of  $\omega_1, \omega_3, \dots, \omega_{n-3}$  that is  $d = \gcd(2n, \binom{2n}{3}, \dots, \binom{2n}{n-3})$ .

In particular, for  $G_s = PGO_8^+$  we obtain

$$\mathfrak{G}_s \simeq \mathbb{Z}[y_1, y_2]/(y_1^2 - 2y_1, y_2^2 - 2y_2, 8y_1, 8y_2, 8y_1y_2).$$

**3.9. Example.** Assume that  $\Lambda/T^* = \langle \sigma \rangle$  has order 3. Then

$$\mathfrak{G}_s \simeq \mathbb{Z}[y]/(y^3 - 3y^2 + 3y, d_1y, d_2y^2),$$

where  $y = (1 - e^\sigma)$  and  $d_1$  (resp.  $d_2$ ) is the greatest common divisor of dimensions of fundamental representations  $\omega_i$ ,  $i = 1 \dots n$  such that  $\bar{\omega}_i = \sigma$  (resp.  $\bar{\omega}_i = 2\sigma$ ).

For the adjoint group of type  $E_6$  we have  $\Lambda/\Lambda_r = \{0, \sigma = \bar{\omega}_1 = \bar{\omega}_5, 2\sigma = \bar{\omega}_2 = \bar{\omega}_6\}$  with  $\bar{\omega}_2 = \bar{\omega}_4 = 0$ . Therefore,  $d_1 = d_2 = \gcd(27, \binom{27}{2}) = 27$ .

**3.10. Example.** Assume that  $\Lambda/T^* = \langle \sigma \rangle$  has order 4. Then

$$\mathfrak{G}_s \simeq \mathbb{Z}[y]/(y^4 - 4y^3 + 6y^2 - 4y, d_1y, d_2y^2, d_3y^3),$$

where  $y = (1 - e^\sigma)$ . For the group  $PGO_{2n}^+$  where  $n$  is odd we have  $\sigma = \bar{\omega}_{n-1}$ ,  $2\sigma = \bar{\omega}_1$  and  $3\sigma = \bar{\omega}_n$ . Therefore,  $d_1 = d_3 = 2^{n-1}$  and  $d_2 = \gcd(\binom{2n}{1}, \binom{2n}{3}, \dots, \binom{2n}{n-2})$ .

#### 4. THE TWISTED $\gamma$ -FILTRATION.

In the present section we introduce and study the twisted  $\gamma$ -filtration.

**4.1.** Let  $\gamma = \ker \epsilon$  denote the augmentation ideal in  $\mathbb{Z}[\Lambda]$ . It is generated by the differences

$$\langle (1 - e^{-\lambda}), \lambda \in \Lambda \rangle.$$

Consider the  $\gamma$ -adic filtration on  $\mathbb{Z}[\Lambda]$

$$\mathbb{Z}[\Lambda] = \gamma^0 \supseteq \gamma \supseteq \gamma^2 \supseteq \dots$$

The  $i$ -th power  $\gamma^i$  is generated by products of at least  $i$  differences.

**4.2. Definition.** We define the filtration on  $K_0(\mathfrak{B}_s)$  (resp. on  $\mathfrak{G}_s$ ) to be the image of the  $\gamma$ -adic filtration on  $\mathbb{Z}[\Lambda]$  via  $\mathfrak{c}$  (resp. via  $q$ ), i.e.

$$\gamma^i K_0(\mathfrak{B}_s) := \mathfrak{c}(\gamma^i) \text{ and } \gamma^i \mathfrak{G}_s := q(\gamma^i K_0(\mathfrak{B}_s)), \quad i \geq 0.$$

So that we have a commutative diagram of surjective group homomorphisms

$$\begin{array}{ccc} \gamma^i & \xrightarrow{\mathfrak{c}} & \gamma^i K_0(\mathfrak{B}_s) \\ & \searrow & \downarrow q \\ & & \gamma^i \mathfrak{G}_s \end{array}$$

**4.3. Lemma.** *The  $\gamma$ -filtration on  $K_0(\mathfrak{B}_s)$  coincides with the filtration introduced in Definition 4.2.*

*Proof.* Since  $K_0(\mathfrak{B}_s)$  is generated by the classes of line bundles,

$$\gamma^i K_0(\mathfrak{B}_s) = \langle c_1([\mathcal{L}_1]) \cdot \dots \cdot c_1([\mathcal{L}_m]) \mid m \geq i, \mathcal{L}_j \in K_0(\mathfrak{B}_s) \rangle.$$

Moreover, each line bundle  $\mathcal{L}$  is the associated bundle  $\mathcal{L} = \mathcal{L}(\lambda)$  for some character  $\lambda \in \Lambda$ . Therefore,  $c_1([\mathcal{L}]) = 1 - [\mathcal{L}^\vee] = \mathfrak{c}(1 - e^{-\lambda})$  (see [3, §2.8]).  $\square$

**4.4. Definition.** Given a  $G_s$ -torsor  $\xi \in H^1(k, G_s)$  and the respective twisted form  $\mathfrak{B} = {}_\xi \mathfrak{B}_s$  we define the twisted filtration on  $\mathfrak{G}_s$  to be the image of the  $\gamma$ -filtration on  $K_0(\mathfrak{B})$  via the composite  $\text{res} \circ q$ , i.e.

$$\gamma_\xi^i \mathfrak{G}_s := q(\text{res}(\gamma^i K_0(\mathfrak{B}))), \quad i \geq 0.$$

Let  $\gamma_\xi^{i/i+1} \mathfrak{G}_s = \gamma_\xi^i \mathfrak{G}_s / \gamma_\xi^{i+1} \mathfrak{G}_s$ . The associated graded ring  $\bigoplus_{i \geq 0} \gamma_\xi^{i/i+1} \mathfrak{G}_s$  will be called the  $\gamma$ -invariant of the torsor  $\xi$  and will be denoted simply as  $\gamma_\xi^*$ .

Note that the Chern classes commute with restrictions, therefore the restriction map  $\text{res}: \gamma^i K_0(\mathfrak{B}) \rightarrow \gamma^i K_0(\mathfrak{B}_s)$  is well-defined. By definition there is a surjective ring homomorphism

$$\gamma^*(\mathfrak{B}) \twoheadrightarrow \gamma_\xi^*.$$

**4.5. Theorem.** *The twisted filtration  $\gamma_\xi^i \mathfrak{G}_s$  can be computed as follows:*

$$\gamma_\xi^i \mathfrak{G}_s = \left\langle \prod_{j=1}^m \binom{\text{ind}(\beta_\xi(\bar{\rho}_{w_j}))}{n_j} (1 - e^{\bar{\rho}_{w_j}})^{n_j} \mid n_1 + \dots + n_m \geq i, w_j \in W \right\rangle.$$

*Proof.* Since the Chern classes commute with restrictions, the image of the restriction  $\text{res}: \gamma^i K_0(\mathfrak{B}) \rightarrow \gamma^i K_0(\mathfrak{B}_s)$  is generated by the products

$$\langle c_{n_1}(\iota_{w_1} g_{w_1}) \cdot \dots \cdot c_{n_m}(\iota_{w_m} g_{w_m}) \mid n_1 + \dots + n_m \geq i, w_1, \dots, w_m \in W \rangle,$$

where  $\{\iota_{w_j}\}$  are the indexes of the respective Tits algebras from 2.5. Applying the Whitney formula for the Chern classes [7, §3.2] we obtain

$$c_j(\iota_w g_w) = \binom{\iota_w}{j} c_1(g_w)^j.$$

Therefore,  $q(\binom{\iota_w}{j} c_1(g_w)^j) = \binom{\iota_w}{j} (1 - e^{-\bar{\rho}_w})^j$ , where  $\iota_w = \text{ind}(\beta_\xi(\bar{\rho}_w))$ .  $\square$

**4.6. Example.** Since  $\gamma^0(X) \simeq \mathbb{Z}$  and  $\gamma^1(X) = \text{Pic}(X)$  is torsion free for every smooth projective  $X$ , we obtain that  $\gamma_\xi^0 \simeq \mathbb{Z}$  and  $\gamma_\xi^1 = 0$  for any  $\xi$ .

**4.7. Example** (Strongly-inner case). If  $\beta_\xi = 0$ , then  $\binom{\iota_{w_j}}{n_j} = 1$  and  $\gamma_\xi^i \mathfrak{G}_s = \gamma^i \mathfrak{G}_s$ .

**4.8. Example** ( $\mathbb{Z}/2\mathbb{Z}$ -case). As in 3.7 assume that  $\Lambda/T^* = \langle \sigma \rangle$  has order 2 and  $\beta_\xi \neq 0$ . Then there is only one non-split Tits algebra  $A = A_{\sigma, \xi}$  and it has exponent 2. Let  $i_A = v_2(\text{ind}(A))$  denote the 2-adic valuation of the index of  $A$ . By definition we have

$$\gamma_\xi^i \mathfrak{G}_s = \langle \binom{2^{i_A}}{n_1} \dots \binom{2^{i_A}}{n_m} 2^{n_1 + \dots + n_m - 1} y \mid n_1 + \dots + n_m \geq i \rangle$$

in  $\mathbb{Z}[y]/(y^2 - 2y, dy)$ , where  $y = 1 - e^\sigma$  and  $d$  is given in 3.7. Observe that modulo the relation  $y^2 = 2y$  these ideals are generated by (for  $j \geq 1$ )

$$\begin{aligned} \gamma_\xi^{2j-1} \mathfrak{G}_s &= \gamma_\xi^{2j} \mathfrak{G}_s = \langle 2^{2j-1} y \rangle & \text{if } i_A = 1; \\ \gamma_\xi^{4j-3} \mathfrak{G}_s &= \gamma_\xi^{4j-2} \mathfrak{G}_s = \langle 2^{4j-2} y \rangle, \quad \gamma_\xi^{4j-1} \mathfrak{G}_s = \gamma_\xi^{4j} \mathfrak{G}_s = \langle 2^{4j-1} y \rangle & \text{if } i_A = 2; \\ \gamma_\xi^1 \mathfrak{G}_s &= \gamma_\xi^2 \mathfrak{G}_s = \langle 2^{i_A} y \rangle, \quad \gamma_\xi^3 \mathfrak{G}_s = \gamma_\xi^4 \mathfrak{G}_s = \langle 2^{i_A+1} y \rangle, \quad \gamma_\xi^5 \mathfrak{G}_s = \langle 2^{i_A+4} y \rangle \dots & \text{if } i_A > 2. \end{aligned}$$

Taking these generators modulo the relation  $dy = 0$  we obtain the following formulas for the second quotient  $\gamma_\xi^2$ :

$$\begin{aligned} \text{if } i_A = 1, \text{ then } \gamma_\xi^2 &= \begin{cases} 0 & \text{if } v_2(d) \leq 1 \\ \mathbb{Z}/2\mathbb{Z} & \text{if } v_2(d) = 2 \\ \mathbb{Z}/4\mathbb{Z} & \text{if } v_2(d) \geq 3 \end{cases} \\ \text{if } i_A > 1, \text{ then } \gamma_\xi^2 &= \begin{cases} 0 & \text{if } v_2(d) \leq i_A \\ \mathbb{Z}/2\mathbb{Z} & \text{if } v_2(d) > i_A \end{cases} \end{aligned}$$

**4.9. Example** ( $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$ -case). As in 3.8 assume that  $\Lambda/T^* = \langle \sigma_1 \rangle \oplus \langle \sigma_2 \rangle$ , where  $\sigma_1, \sigma_2$  have order 2. This is the case for the adjoint group  $PGO_{2n}^+$  where  $n$  is even. Assume that  $n = 4$  which corresponds to the group of type  $D_4$ , i.e.  $PGO_8^+$ . Let  $C^+$  and  $C^-$  denote the Tits algebras corresponding to the generators  $\sigma_1 = \bar{\omega}_3$  and  $\sigma_2 = \bar{\omega}_4$ . Let  $A$  denote the Tits algebra corresponding to the sum  $\sigma_1 + \sigma_2$ . Note that  $C^+ \times C^-$  is the even part of the Clifford algebra of the algebra with involution  $A$  and  $[A] = [C^+ \otimes C^-]$  in  $Br(k)$ .

By definition we have in  $\mathbb{Z}[y_1, y_2]$

$$\gamma_\xi^i \mathfrak{G}_s = \langle \binom{ind C_+}{n_1} y_1^{n_1} \cdot \binom{ind C_-}{n_2} y_2^{n_2} \cdot \binom{ind A}{n_3} (y_1 + y_2 - y_1 y_2)^{n_3} \mid n_1 + n_2 + n_3 \geq i \rangle.$$

Modulo the relations  $(y_1^2 - 2y_1, y_2^2 - 2y_2, 8y_1, 8y_2, 8y_1 y_2)$  we obtain that

$$\gamma_\xi^2 \mathfrak{G}_s \simeq \frac{(ind C_+) \mathbb{Z}}{8\mathbb{Z}} \oplus \frac{(ind C_-) \mathbb{Z}}{8\mathbb{Z}} \oplus \frac{(ind A) \mathbb{Z}}{8\mathbb{Z}}$$

## 5. TORSION IN THE $\gamma$ -FILTRATION.

In the present section we show how the twisted  $\gamma$ -filtration can be used to construct nontrivial torsion elements in the  $\gamma$ -ring of the twisted form  $\mathfrak{B}$  of a variety of Borel subgroups.

5.1. For simplicity we consider only the case of  $G_s$  (see Examples 3.7 and 4.8) with  $\Lambda/T^* = \langle \sigma \rangle$  of order 2. Let  $d$  denote the g.c.d. of dimensions of fundamental representations corresponding to  $\sigma$ .

Given a  $G_s$ -torsor  $\xi \in H^1(k, G_s)$  let  $i_A$  denote the 2-adic valuation of the index of the Tits algebra  $A = A_{\sigma, \xi}$ . Let  $\mathfrak{B} = {}_\xi \mathfrak{B}_s$  denote the twisted form of the variety of Borel subgroups of  $G_s$  by means of  $\xi$ . Consider the respective twisted filtration  $\gamma_\xi^i \mathfrak{G}_s$  on  $\mathfrak{G}_s$ .

5.2. **Proposition.** *Assume that  $v_2(d) > i_A \geq 3$ . Then for each  $\lambda \in \Lambda$  such that  $\bar{\lambda} = \sigma$  there exists a non-trivial torsion element of order 2 in  $\gamma^2(\mathfrak{B})$ . Moreover, its image in  $\gamma_\xi^2 = \mathbb{Z}/2$  (via  $q$ ) is non-trivial and in  $\gamma^2(\mathfrak{B}_s)$  (via  $res$ ) is trivial.*

*Proof.* The proof of this result was inspired by the proof of [8, Prop.4.13].

Let  $g = [\mathcal{L}(\lambda)]$  denote the class of the associated line bundle. Using the formula for the first Chern class of a tensor product of line bundles for  $K_0$  we obtain

$$c_1(g)^2 = 2c_1(g) - c_1(g^2).$$

Hence,

$$c_1(g)^4 = (2c_1(g) - c_1(g^2))^2 = 4c_1(g)^2 - 4c_1(g)c_1(g^2) + c_1(g^2)^2.$$

Therefore,

$$\eta = 4c_1(g)^3 - c_1(g)^4 = 4c_1(g)^2 - c_1(g^2)^2 \in \gamma^3 K_0(\mathfrak{B}_s).$$

We claim that the class of  $2^{i_A-3}\eta$  gives the desired torsion element.

Indeed,  $c_1(g^2) = c_1([\mathcal{L}(2\lambda)])$ . Since  $2\lambda \in T^*$ ,  $[\mathcal{L}(2\lambda)] \in \mathfrak{c}(T^*)$  and, therefore, by [5, Cor.3.1]  $c_1(g^2) \in \gamma^1 K_0(\mathfrak{B})$ . Moreover, we have  $2^{i_A-1}c_1(g)^2 = c_2(2^{i_A}g)$ , where  $2^{i_A}g \in K_0(\mathfrak{B})$ . Hence,  $2^{i_A-1}c_1(g)^2 \in \gamma^2 K_0(\mathfrak{B})$ . Combining these together we obtain that  $2^{i_A-3}\eta \in \gamma^2 K_0(\mathfrak{B})$ .

Now since  $2^{i_A-3}\eta \in \gamma^2 K_0(\mathfrak{B})$  its image in  $\gamma_\xi^2 \mathfrak{G}_s$  can be computed as

$$q(2^{i_A-3}\eta) = 2^{i_A-3}q(\eta) = 2^{i_A-1}q(c_1(g)^2) = 2^{i_A-1}(1 - e^{-\sigma})^2 = 2^{i_A}y.$$

But  $q(2^{i_A-3}\eta) \notin \gamma_\xi^3 \mathfrak{G}_s = \langle 2^{i_A+1}y \rangle$ . Therefore,  $2^{i_A-3}\eta \notin \gamma^3 K_0(\mathfrak{B})$ .

From the other hand side  $2^{i_A-2}\eta = 2^{i_A}c_1(g)^3 + 2^{i_A-2}c_1(g)^4$  is in  $\gamma^3 K_0(\mathfrak{B})$ . So the class of  $2^{i_A-3}\eta$  gives the desired torsion element of order 2.  $\square$

5.3. **Example.** Let  $G_s = HSpin_{2n}$  be a half-spin group of rank  $n \geq 4$ . So  $G_s$  is of type  $D_n$ , where  $n$  is even,  $\Lambda/T^* = \langle \sigma = \bar{\omega}_1 \rangle \simeq \mathbb{Z}/2\mathbb{Z}$  and according to Example 3.7 we have  $d = 2^{v_2(n)+1}$ . Let  $\xi \in H^1(k, G_s)$  be a non-trivial torsor. Then there is only one Tits algebra  $A = A_{\sigma, \xi}$ ; it has exponent 2 and index  $2^{i_A}$  such that  $i_A \leq v_2(n) + 1$ .



Recall that each such torsor corresponds to an algebra with orthogonal involution  $(A, \delta)$  with trivial discriminant and trivial component of the Clifford algebra. The respective twisted form  $\mathfrak{B} = {}_\xi \mathfrak{B}_s$  then corresponds to the variety of Borel subgroups of the group  $PGO^+(A, \delta)$ .

Applying the proposition to this case we obtain that for any such algebra  $(A, \delta)$  where  $8 \mid \text{ind}(A)$  and  $A$  is non-division, there exists a non-trivial torsion element of order 2 in  $\gamma^2(\mathfrak{B})$  that vanishes over a splitting field of  $(A, \delta)$ .

**5.4. Lemma.** *The  $\gamma$ -filtration on  $K_0(\mathfrak{B}_s)$  is generated by the first Chern classes  $c_1([\mathcal{L}(\omega_i)])$ ,  $i = 1 \dots n$ , i.e.*

$$\gamma^i K_0(\mathfrak{B}_s) = \langle \prod_{j \in 1 \dots n} c_1([\mathcal{L}(\omega_j)]) \mid \text{the number of elements in the product} \geq i \rangle.$$

*In particular, the second quotient  $\gamma^2(\mathfrak{B}_s)$  is additively generated by the products*

$$\gamma^2(\mathfrak{B}_s) = \langle c_1([\mathcal{L}(\omega_i)])c_1([\mathcal{L}(\omega_j)]) \mid i, j \in 1 \dots n \rangle.$$

*Proof.* Each  $b \in K_0(\mathfrak{B}_s)$  can be written as a linear combination  $b = \sum_{w \in W} a_w g_w$ . Therefore, any Chern class of  $b$  can be expressed in terms of  $c_1(g_w)$ .

Each  $\rho_w$  can be written uniquely as a linear combination of fundamental weights  $\{\omega_1, \dots, \omega_n\}$ . Therefore, by the formula for the Chern class of the tensor product of line bundles [2, 8.2], each  $c_1(g_w)$  can be expressed in terms of  $c_1([\mathcal{L}(\omega_i)])$ .  $\square$

**5.5. Example.** Let  $G_s$  be an adjoint group of type  $E_7$  and let  $\xi \in H^1(k, G_s)$  be a non-trivial  $G_s$ -torsor. Then there is only one nonsplit Tits algebra  $A = A_{\sigma, \xi}$  of exponent 2 and  $i_A \leq 3$ . Let  $\mathfrak{B} = {}_\xi \mathfrak{B}_s$  be the respective twisted flag variety.

By Lemma 5.4 any element of  $\gamma^2(\mathfrak{B})$  can be written as

$$x = \sum_{ij} a_{ij} c_1([\mathcal{L}(\omega_i)])c_1([\mathcal{L}(\omega_j)]) \in \gamma^2(\mathfrak{B})$$

for certain coefficients  $a_{ij} \in \mathbb{Z}$ . Since  $\sigma = \bar{\omega}_7 = \bar{\omega}_5 = \bar{\omega}_2$  and  $\bar{\omega}_1 = \bar{\omega}_3 = \bar{\omega}_4 = \bar{\omega}_6 = 0$ , we obtain that

$$q(x) = C \cdot 2y \in \gamma_\xi^2, \text{ where } C = a_{25} + a_{27} + a_{57} + a_{22} + a_{55} + a_{77}.$$

Therefore,  $q(x) \neq 0$  in  $\gamma_\xi^2$  if and only if  $4 \nmid C$  and  $i_A \leq 2$ .

Consider the class  $\mathfrak{c}(\theta) \in \gamma^2 K_0(\mathfrak{B}_s)$  of the special cycle  $\theta$  constructed in [4, Def.3.3]. Note that the image of  $\theta$  in  $CH^2(\mathfrak{B})$  can be viewed as a generalization of the Rost invariant for split adjoint groups (see Remark 5.7).

If  $i_A = 1$ , then by [4, Prop.6.5] we know that  $\mathfrak{c}(\theta) \in \gamma^2(\mathfrak{B})$  is a non-trivial torsion element. If  $i_A = 2$ , then following the proof of [4, Prop.6.5] we obtain that  $2\mathfrak{c}(\theta) \in \gamma^2(\mathfrak{B})$ .

We claim that if  $i_A \leq 2$ , then  $x = 2\mathfrak{c}(\theta)$  is non-trivial. Indeed, in this case  $4 \nmid C = a_{22} + a_{55} + a_{77} = 6$ , therefore, we have  $q(x) \neq 0$ , and  $x \neq 0$  in  $\gamma^2(\mathfrak{B})$ . In particular, this shows that for  $i_A = 1$  the order of the special cycle  $\theta$  in  $\gamma^2(\mathfrak{B})$  is divisible by 4.

**5.6. Example.** Let  $\xi \in H^1(k, PGO_8^+)$ . Applying the same arguments as in 5.5 to Example 4.9 we obtain that if  $\text{ind}(A), \text{ind}(C_+), \text{ind}(C_-) \leq 4$ , then  $2\mathfrak{c}(\theta) \in \gamma^2(\mathfrak{B})$  is non-trivial.

We finish by the following remark that provides another motivation for the study of the torsion part of  $\gamma^*(\mathfrak{B})$

**5.7. Remark.** Recall that by the Riemann-Roch theorem the second Chern class induces a surjection  $c_2: Tors \gamma^2(\mathfrak{B}) \twoheadrightarrow Tors CH^2(\mathfrak{B})$  [8, Cor.2.15], where the latter group is isomorphic to the cohomology quotient [12, Thm.2.1]

$$\frac{\ker(H^3(k, \mathbb{Q}/\mathbb{Z}(2)) \rightarrow H^3(k(\mathfrak{B}), \mathbb{Q}/\mathbb{Z}(2)))}{\oplus_{\chi \in \Lambda/\Lambda_r} N_{k_\chi/k}(k_\chi^* \cup \beta_\xi(\chi))},$$

where  $k_\chi$  denotes the fixed subfield of  $\chi$ . Therefore, the group  $Tors \gamma^2(\mathfrak{B})$  can be viewed as an upper bound for the group of cohomological invariants of  $G_s$ -torsors in degree 3.

**Acknowledgments.** I am grateful to Alexander Merkurjev for pointing out Remark 3.2. This work was supported by NSERC Discovery 385795-2010 and Accelerator Supplement 396100-2010 grants.

#### REFERENCES

1. Bourbaki, N. Éléments de Mathématique, Groupes et Algèbres de Lie 7-8 et 9, 1982.
2. Calmès, B., Petrov, V., Zainoulline, K. Invariants, torsion indices and oriented cohomology of complete flags. Preprint arXiv 2010, 36pp.
3. Demazure, M. Désingularisation des variétés de Schubert généralisées. Ann. Sci. École Norm. Sup. (4) 7 (1974), 53–88.
4. Garibaldi, S., Zainoulline, K. The  $\gamma$ -filtration and the Rost invariant. Preprint arXiv 2010, 19pp.
5. Gille, S., Zainoulline, K. Equivariant pretheories and invariants of torsors, Preprint arXiv 2010, 8pp.
6. Grothendieck, A. Théorie des Intersections et Théorème de Riemann-Roch (SGA 6). Exposé 0, LNM 225, Springer-Verlag, 1971.
7. Fulton, W. Intersection theory. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge., 2. Springer-Verlag, Berlin, 1998. xiv+470 pp.
8. Karpenko, N. Codimension 2 cycles on Severi-Brauer varieties. K-Theory 13 (1998), no. 4, 305–330.
9. Knus, M.-A., Merkurjev, A., Rost, M., Tignol, J.-P. The Book of Involutions. American Mathematical Society Colloquium Publications, 44. American Mathematical Society, Providence, RI, 1998.
10. Merkurjev A. Equivariant K-theory. J. Handbook of K-theory. Vol. 1, 2, Springer, Berlin, 925-954, 2005.
11. Panin, I. On the Algebraic K-Theory of Twisted Flag Varieties. K-Theory J. 8 (1994), 541–585.
12. Peyre, E. Galois cohomology in degree three and homogeneous varieties. K-theory 15 (1998), 99–145.
13. Steinberg, R., On a Theorem of Pittie, Topology Vol. 14 (1975), 173–177.
14. Tits, J. Représentations linéaires irréductibles d’un groupe réductif sur un corps quelconque. (French) J. Reine Angew. Math. 247 (1971), 196–220